



Incidence of Type 1 Diabetes in Children and Adolescents During the COVID-19 Pandemic in Germany: Results From the DPV Registry

<https://doi.org/10.2337/dc21-0969>

Clemens Kamrath¹,
Joachim Rosenbauer,^{2,3}
Alexander J. Eckert,^{3,4} Kai Siedler,⁵
Heike Bartelt,⁶ Daniela Klose,⁷
Marina Sindichakis,⁸ Silke Herrlinger,⁹
Valentina Lahn,¹⁰ and Reinhard W. Holl^{3,4}

OBJECTIVE

The aim of this study was to investigate the incidence of type 1 diabetes in children and adolescents during the coronavirus disease 2019 (COVID-19) pandemic in Germany compared with previous years.

RESEARCH DESIGN AND METHODS

Based on data from the multicenter German Diabetes Prospective Follow-up Registry, we analyzed the incidence of type 1 diabetes per 100,000 patient-years in children and adolescents from 1 January 2020 through 30 June 2021. Using Poisson regression models, expected incidences for 2020/21 were estimated based on the data from 2011 to 2019 and compared with observed incidences in 2020/21 by estimating incidence rate ratios (IRRs) with 95% CIs.

RESULTS

From 1 January 2020 to 30 June 2021, 5,162 children and adolescents with new-onset type 1 diabetes in Germany were registered. The observed incidence in 2020/21 was significantly higher than the expected incidence (24.4 [95% CI 23.6–25.2] vs. 21.2 [20.5–21.9]; IRR 1.15 [1.10–1.20]; $P < 0.001$). IRRs were significantly elevated in June 2020 (IRR 1.43 [1.07–1.90]; $P = 0.003$), July 2020 (IRR 1.48 [1.12–1.96]; $P < 0.001$), March 2021 (IRR 1.29 [1.01–1.65]; $P = 0.028$), and June 2021 (IRR 1.39 [1.04–1.85]; $P = 0.010$).

CONCLUSIONS

A significant increase in the incidence of type 1 diabetes in children was observed during the COVID-19 pandemic, with a delay in the peak incidence of type 1 diabetes by ~3 months after the peak COVID-19 incidence and also after pandemic containment measures. The underlying causes are yet unknown. However, indirect rather than direct effects of the pandemic are more likely to be the cause.

The rate of new cases of coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) increased remarkably during March 2020 and peaked in early April 2020 in Germany (1,2). In response to the pandemic, nationwide health policy measures to achieve social distancing, such as restrictions on social contacts, school closures, and the general recommendation

¹Center of Child and Adolescent Medicine, Justus Liebig University, Giessen, Germany

²Institute for Biometrics and Epidemiology, German Diabetes Center, Leibniz Center for Diabetes Research, Heinrich Heine University Dusseldorf, Dusseldorf, Germany

³German Center for Diabetes Research, Munich-Neuherberg, Germany

⁴Institute of Epidemiology and Medical Biometry, Central Institute of Biomedical Technology, Ulm University, Ulm, Germany

⁵Hospital for Children and Adolescents, Helios Clinics Pforzheim, Pforzheim, Germany

⁶Hospital for Children and Adolescents, University of Leipzig, Leipzig, Germany

⁷University Children's Hospital Heidelberg, Heidelberg, Germany

⁸Hospital for Children and Adolescents, Klinikum Traunstein, Traunstein, Germany

⁹Professor Hess Parent-Child Center, Bremen Central Clinic, Bremen, Germany

¹⁰Altonaer Kinderkrankenhaus, Hamburg, Germany

Corresponding author: Clemens Kamrath, clemens.kamrath@paediat.med.uni-giessen.de

Received 6 May 2021 and accepted 16 December 2021

This article contains supplementary material online at <https://doi.org/10.2337/figshare.17282138>.

© 2022 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

to stay at home to contain the spread of SARS-CoV-2, were implemented. Thereafter, the incidence of COVID-19 in Germany stabilized at a lower level from June 2020 to September 2020. In early October 2020, however, COVID-19 cases showed another sharp increase in a second wave, with a peak in late December 2020, which was followed by a third wave starting in March 2021 and lasting until June 2021 (1,2).

Type 1 diabetes is a chronic autoimmune disease that is influenced by both genetic and environmental factors (3). The incidence of type 1 diabetes increased by an average annual rate of 3–4% over the past three decades (4). There is evidence for a role of infections in the pathogenesis of type 1 diabetes (5,6). Common respiratory infections in early childhood have been shown to be a risk factor for the development of type 1 diabetes (7–9). Infection with SARS-CoV-2 was suspected to lead to the development of new-onset diabetes in children (10). However, it is not known whether COVID-19 is associated with an increase in the incidence of type 1 diabetes (11,12). On the other hand, efforts to control the COVID-19 pandemic, such as social isolation, led to a marked decrease in common infections in children (13,14), and this could also affect the incidence of type 1 diabetes. It is therefore essential to clarify whether the incidence of type 1 diabetes changed during this pandemic. We have previously shown that there was no increase in the incidence in type 1 diabetes during the first wave of the COVID-19 pandemic in Germany from mid-March to mid-May 2020 (15).

The aim of this study was to extend this analysis with longer follow-up. We therefore investigated the frequency of new-onset type 1 diabetes in children and adolescents during the COVID-19 pandemic in 2020 and the first half of 2021 compared with the years 2011 to 2019 in Germany using a large population-based clinical database.

RESEARCH DESIGN AND METHODS

Data Source and Study Population

This study used data from the DPV (German Diabetes Prospective Follow-up Registry; i.e., Diabetes-Patienten-Verlaufsdokumentation) of children and adolescents living in Germany and aged between 6 months and <18 years at the diagnosis of new-onset type 1 diabetes

from 1 January 2020 through 30 June 2021. The control group consisted of 22,987 children and adolescents with new-onset type 1 diabetes in the same age range diagnosed from 2011 to 2019 in Germany. The DPV has a nationwide coverage of >90% of pediatric patients with type 1 diabetes in Germany and comprises 257 pediatric diabetes centers in Germany (16). Twice a year, locally collected longitudinal data are pseudonymized and transmitted for central plausibility checks and analyses to Ulm University, Ulm, Germany. Inconsistent data are reported back to participating centers for validation and/or correction. Data are then completely anonymized for analysis. Verbal or written informed consent for participation in the DPV was obtained from patients or their guardians. The ethics committee of Ulm University in Ulm, Germany, approved the analysis of anonymized data from the DPV. This study followed the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) reporting guideline for cohort studies.

The severity of the COVID-19 pandemic was measured as the weekly incidence rates of new COVID-19 cases per 100,000 population. In accordance with the international standards of the World Health Organization, all laboratory confirmations of SARS-CoV-2, irrespective of the presence or severity of clinical symptoms, were considered as COVID-19 cases. The incidence of confirmed COVID-19 cases was derived from the official statistics of the Robert Koch Institute, Berlin, Germany (17); the population data were taken from the German Federal Statistical Office (18).

The Stringency index, which is a composite index containing nine indicators of containment and closure policies, originated from the Oxford Covid-19 Government Response Tracker (OxCGRT) (19).

Variables

Demographic data included year, month (and additionally week for 2018 to 2020/21 data), and age at diabetes onset, sex, and immigrant background (patient or at least one parent born outside of Germany). Clinical data from diabetes onset included BMI (calculated as weight in kilograms divided by height in meters squared), HbA_{1c} (% [mmol/mol]), and presence of autoantibodies.

BMI values were transformed to SD scores based on German reference values (KiGGS [German Health Interview and Examination Survey for Children and Adolescents]) by applying the Box-Cox transformation method (20). In order to adjust for different laboratory methods, local HbA_{1c} values were mathematically standardized to the DCCT (Diabetes Control and Complications Trial) reference range (4.05–6.05%) using the multiple of the mean transformation method. Autoantibody measurements included autoantibodies against islet cells, glutamic acid decarboxylase, tyrosine phosphatase, insulin, and zinc transporter 8. Positive autoantibody status was defined as any measure of these autoantibodies above the upper limit of normal, according to the reference values of the respective laboratory of the treating center, at the diagnosis of diabetes.

We analyzed the incidence of type 1 diabetes for the whole period from 1 January 2020 to 30 June 2021, separately for the year 2020 and the first half of the year 2021, as well as for five different periods related to the activity of the COVID-19 pandemic: the prepandemic period, consisting of January and February 2020; the first wave of the pandemic, consisting of March to May 2020; a stable phase with a relatively low rate of new infections from June to September 2020; the second wave from October 2020 to February 2021; and the third wave from March 2021 to the end of June 2021 (1). Furthermore, we examined the incidence of type 1 diabetes by month from January 2020 to June 2021.

Statistical Analysis

Medians and interquartile ranges (IQRs) are provided for the description of continuous variables, and frequencies and percentages are used for the description of categorical variables. For each timeframe analyzed (total period from January 2020 to June 2021, calendar years, the first halves of calendar years, COVID-19–related periods, or months), we related the number of new cases to the respective number of person-years at risk to estimate the incidence per 100,000 person-years with 95% CIs.

For the analysis of the incidence by calendar year, we applied a multivariable Poisson trend regression model

(model 1) to incidence data from 2011 to 2019 that included year at diabetes onset (as continuous term), age group at diabetes onset (6 months to <6 years, 6 to <12 years, and 12 to <18 years), and sex as independent variables. Based on this trend model, we estimated the expected incidence for 2011 to 2020/21, standardized for age group and sex, with the sum of person-years of each year between 2011 and 2020 as reference distribution.

We used a respective model with year at diabetes onset included as a categorical independent term (model 2) to estimate incidences for 2011 to 2019, standardized for age group and sex, using the same reference distribution of person-years at risk. Furthermore, we applied a multivariable Poisson regression model that included age group and sex as independent variables (model 3) to adjust the observed incidence in 2020/21 to the same reference distributions of the person-years at risk. Finally, a Poisson regression model that included a binary variable indicating observed and predicted data as independent variables (model 4) was used to compare the observed with the expected incidence of type 1 diabetes. The ratio between the observed and expected incidence of type 1 diabetes was presented as adjusted incidence rate ratio (IRR), with the corresponding 95% CIs and two-sided *P* values from the Wald test. CIs of observed and expected incidences in strata were adjusted for multiple inferences according to the Bonferroni procedure. CIs for estimated period/month-specific IRRs were adjusted for multiple inferences according to the Bonferroni procedure, and corresponding *P* values were adjusted for multiple testing using the Bonferroni-Holm method.

Furthermore, we analyzed the incidence data stratified by age group and sex, applying the same statistical approach (with necessary adaptations).

To analyze incidence data by period (January to February 2020, March to May 2020, June to September 2020, October 2020 to February 2021, and March to June 2021) or month (January 2020 to June 2021), the Poisson regression models described before were extended appropriately. For period-specific or month-specific analyses, models 1 and 2 were extended to include the period or month of diabetes onset and

a term for an interaction of period or month at diabetes onset by year at diabetes onset. Model 3 was extended to include only period or month of diabetes onset. Model 4 additionally included a term for interaction of the binary variable indicating observed and predicted data and period or month.

Furthermore, rates of autoantibody negativity in patients with newly diagnosed type 1 diabetes in 2020/21 were compared with those in 2018 and 2019 via multivariable log-binomial regression, adjusted for age group, sex, and immigrant background and additionally for number of measured antibodies in cases of autoantibody negativity. The analysis of autoantibody negativity was performed for the whole period from January 2020 to June 2021 versus 2018/2019, total calendar years, for the first halves of the years, and stratified by periods based on the total onset cohorts.

Differences in frequencies of autoantibody negativity were presented as adjusted absolute percentage differences with the corresponding 95% CIs.

A two-sided *P* value <0.05 was considered statistically significant. All analyses were performed with SAS version 9.4 (SAS Institute, Inc., Cary, NC).

Data and Resource Availability

Access to the data is possible by remote data processing upon request and approval from the DPV board.

RESULTS

Study Cohort

In the period from 1 January 2020 to 30 June 2021, the DPV registered 5,162 children and adolescents (55.8% male) with new-onset type 1 diabetes from 236 diabetes centers in Germany, 3,338 patients from 2020 and 1,824 patients from the first half of 2021. The median age of the cohort was 9.7 years (IQR 5.9–12.8). The median HbA_{1c} level was 11.4% (IQR 9.9–13.1 [101.3 mmol/mol (84.8–119.6)]).

Table 1 provides a descriptive overview of the 2020/21 study population and a comparison with the 2018–2019 cohort. We compared the weekly numbers of cases of children and adolescents with new-onset type 1 diabetes in 2020 and the first half of 2021 with those of the 2 previous years. Figure 1 shows the weekly cumulative case

numbers for each year. An increase in the case numbers in the years 2020 and 2021 compared with the years 2019 and 2018 is evident onward from calendar week 20 in 2020 and week 11 in 2021, respectively.

Annual Change in Incidence of Type 1 Diabetes From 2011 to 2019

From 2011 to 2019, the incidence of type 1 diabetes increased by 2.4% per year (95% CI 1.9–2.9; *P* < 0.001) (Fig. 2). This increase was higher in male than in female participants (male, 2.7% per year [95% CI 2.0–3.4]; *P* < 0.001 and female, 2.0% per year [95% CI 1.2–2.7]; *P* < 0.001). The increase in the incidence of type 1 diabetes between 2011 and 2019 was most pronounced among adolescents aged 12–17 years (4.3% per year [95% CI, 3.4–5.2]; *P* < 0.001), lower in those aged 6–11 years (1.7% per year [95% CI 0.9–2.5]; *P* < 0.001), and lowest in children aged <6 years (1.1% per year [95% CI 0.1 to 2.1]; *P* = 0.033).

Observed Compared With Expected Incidence of Type 1 Diabetes in 2020/21

For the period from 1 January 2020 through 30 June 2021, the observed incidence was 24.4 (95% CI 23.6–25.2). This observed incidence was significantly higher than the expected incidence of 21.2 (95% CI 20.5–21.9); the corresponding IRR was 1.15 (95% CI 1.10–1.20; *P* < 0.001) (Table 2). The IRR was similar for both sexes (female, 1.14 [95% CI 1.07–1.21]; *P* < 0.001 and male, 1.16 [95% CI 1.10–1.23]; *P* < 0.001) (Table 2).

The increase in the observed compared with the expected incidence in 2020/21 was significant in the age groups of children aged <6 years (IRR 1.23 [95% CI 1.13–1.33]; *P* < 0.001) and 6–11 years (IRR 1.18 [95% CI 1.11–1.26]; *P* < 0.001), while no significant increase could be detected in adolescents aged 12–17 years (IRR 1.06 [95% CI 0.98–1.13]; *P* = 0.13) (Table 2).

Separately for 2020 and for the first half of 2021, the observed incidences (23.7 [95% CI 22.8–24.6] and 25.8 [24.4–27.2], respectively) were each significantly higher than the expected incidences (20.9 [20.0–21.8] and 21.8 [20.5–23.1], respectively), with IRRs of 1.13 (1.08–1.19; *P* < 0.001) and 1.18

Table 1—Characteristics of patients with new-onset type 1 diabetes from 2020 and first half of 2021 and from 2018 and 2019

Variable	2020/21 (n = 5,162)	2018 (n = 2,740)	2019 (n = 2,903)
Median age at diabetes diagnosis (IQR), years	9.7 (5.9–12.8)	9.8 (6.0–13.1)	9.7 (5.9–12.9)
Male sex, %	55.8	55.0	54.9
Median HbA _{1c} at diabetes diagnosis (IQR), % [mmol/mol]	11.4 (9.9–13.1) [101.3 (84.8–119.6)]	11.0 (9.5–12.7) [97.2 (80.7–115.6)]	11.2 (9.7–12.9) [98.3 (82.7–117.4)]
Median BMI at diabetes diagnosis (IQR), SD score	−0.25 (−1.10 to 0.56)	−0.31 (−1.11 to 0.53)	−0.24 (−1.10 to 0.59)
Immigrant background, %	26.0	25.8	27.3

For the demonstration of the demographic and clinical data, the cohort of 2020/21 was compared with children and adolescents aged between 6 months and <18 years at the diagnosis of type 1 diabetes in the 2 previous years 2019 and 2018 in Germany.

(1.10–1.27; $P < 0.001$), respectively (Fig. 2 and Table 2).

Observed Compared With Expected Incidence of Type 1 Diabetes by Different Pandemic-Related Periods or Months in 2020/21

An analysis of the observed and expected incidences of type 1 diabetes in the five different periods related to the activity of the COVID-19 pandemic revealed a significantly higher observed than expected incidence in summer 2020 from June to September 2020 (IRR 1.27 [95% CI 1.13–1.43]; $P < 0.001$) and in spring 2021 from March to June 2021 (IRR 1.27 [1.13–1.42]; $P < 0.001$) (Fig. 3 and Supplementary Fig. 1). Stratified by age group, we found a significantly higher observed

than expected incidence from June to September 2020 and from March to June 2021 for the children aged <12 years, while for adolescents aged 12–17 years, the observed incidence was only higher than the expected in the period from June to September 2020 (Table 2).

Figure 4 shows the monthly IRRs with the corresponding 95% CIs. Compared with the expected monthly incidence, the observed incidence was significantly higher in June 2020 (IRR 1.43 [95% CI 1.07–1.90]; $P = 0.003$), July 2020 (IRR 1.48 [1.12–1.96]; $P < 0.001$), March 2021 (IRR 1.29 [1.01–1.65]; $P = 0.028$), and June 2021 (IRR 1.39 [1.04–1.85]; $P = 0.010$). Supplementary Table 1 provides the observed and expected incidences by month from January 2020 to

June 2021 and the corresponding IRRs. Compared with the 7-day incidence of COVID-19 cases, COVID-19–related deaths, and the Stringency index as a measure of the pandemic containment measures presented in Supplementary Fig. 2, the increase in the incidence of type 1 diabetes followed the peak incidences of the pandemic and the countermeasures taken to control it by ~3 months.

Autoantibody Negativity in 2020/21 Compared With 2018–2019

Data of autoantibody measurements were available in 3,851 (74.6%) of 5,162 patients in 2020 and the first half of 2021 and in 4,205 (74.5%) of 5,643 patients in 2018 and 2019. For the period from 1 January 2020 through 30 June 2021, the adjusted rates of autoantibody negativity did not differ from 2018 and 2019 (2020/21 vs. 2018/19, 6.1% [95% CI 5.3–7.0] vs. 6.4% [5.7–7.3]; absolute difference 2020/21 vs. 2018/19, −0.3% [−1.1 to 0.6]; $P = 0.47$). In addition, separately for the year 2020 and for the first half of 2021, the adjusted rates of autoantibody negativity did not differ from 2018 and 2019 (difference 2020 vs. 2018/19, −0.5% [−1.5 to 0.5]; $P = 0.32$ and difference first halves of 2021 vs. 2018/19, −0.2% [−1.5 to 1.2]; $P = 0.83$).

Moreover, during both periods with an increased incidence of type 1 diabetes, there was also no significant difference for the adjusted rate of autoantibody negativity (absolute difference June to September 2020 vs. June to September 2018/19, 0.7% [95% CI −1.0 to 2.4]; $P = 0.40$ and absolute difference February to June 2021 vs. February to June 2018/19, −0.1% [−1.7 to 1.6]; $P = 0.95$).




Figure 1—Cumulative number of new cases of type 1 diabetes in children and adolescents in Germany over the years 2018 (black), 2019 (blue), and 2020 (red) and the first half of 2021 (green). Week-specific information on the time of diabetes onset was missing for seven patients in 2021, 12 patients in 2020, and nine patients in 2019 and 16 patients in 2018. These patients were not included in the illustration.




Figure 2—Incidence of type 1 diabetes in children in the year 2020 (A) and the first half of 2020 and 2021 (B) compared with the years 2011 to 2019. Based on a multivariable Poisson trend regression model on data from 2011 to 2019 that included year at diabetes onset (as continuous term), age group at diabetes onset, and sex as independent variables, the expected incidence (as rate per 100,000 person-years) for 2011 to 2020, as well for the first half of the years 2011 to 2021, standardized for age group and sex, with the sum of person-years of each year between 2011 and 2020/21 as reference distribution, was estimated (dark blue line with corresponding 95% CI [light blue area]). A multivariable Poisson regression model with year at diabetes onset included as categorical independent term was used to estimate incidences for 2011 to 2019 standardized for age group and sex using the same reference distribution of person-years at risk (red circles). Furthermore, a multivariable Poisson regression model that included age group and sex as independent variables was applied to adjust the observed incidence in 2020 and the first half in 2021 to the same reference distributions of the person-years at risk (red circles). The vertical bars indicate the corresponding 95% CIs of the estimated observed incidences. The number of children registered in DPV with new-onset type 1 diabetes is presented for every year from 2011 to 2020 (A) and for every first half of the years 2011 to 2021 (B).

CONCLUSIONS

This population-based study found an increase in the observed versus expected incidence of type 1 diabetes in children and adolescents during the COVID-19 pandemic. The peak incidence of type 1 diabetes followed the peak incidence of

COVID-19 and also the pandemic containment measures by ~ 3 months. Except for the increase in March 2021, the increase in the incidence of type 1 diabetes was mainly due to the lack of a seasonal decline during the summer months.

Reports from diabetes registries in Norway (21) and Austria (22) indicate that the incidence of type 1 diabetes in children aged <15 years stopped rising from 2004 to 2012 and from 2012 to 2017, respectively, especially in young children. In contrast, our study shows a preserved increase in the incidence of type 1 diabetes in childhood between 2011 and 2019. Consistent with the studies from Norway and Austria mentioned above, the increase declined with decreasing age at diabetes onset. In contrast to this trend in recent years, the most pronounced increase of incidence of type 1 diabetes during the COVID-19 pandemic was observed in this youngest age group.

The results of this study are different from those presented in a previous short communication from our group, demonstrating no increase in type 1 diabetes incidence during the first wave of the COVID-19 pandemic (15). However, this is not a contradiction. The previous observation interval was limited to the first 2 months of the pandemic and the lockdown from mid-March to mid-May (calendar week 20). As this extension of the observation period shows, the incidence increased just after the end of the previous observation period. This also demonstrates the importance of monitoring the course of the pandemic in the longer term.

Our study is consistent with the results of a smaller study from Finland (23). In this study, children with new-onset type 1 diabetes from 1 April to 31 October from 2016 to 2020 in the Helsinki University Hospital were analyzed. During the pandemic period, 84 children with newly diagnosed type 1 diabetes were registered, as compared with 53–62 children each year in the prepandemic periods of 2016 to 2019. As in our study, the most significant increase in the number of cases occurred in the 2 summer months of June and July 2020. Importantly, all 33 children diagnosed in 2020 who were tested for SARS-CoV-2 antibodies were negative. In contrast to this report and our results, a study involving 13 pediatric diabetes centers in Lombardy, Italy, found no significant increase in the incidence of type 1 diabetes in 2020 compared with 2019 (24). The reported new case number of 256 in 2020 was not significantly

Table 2—Observed versus expected incidence of type 1 diabetes (per 100,000 person-years) during year 2020 and first half of 2021, as well as during both periods with increased observed incidence

	Observed incidence in 2020/21 (95% CI)	Expected incidence for 2020/21 based on data from 2011–2019 (95% CI)	IRR for observed vs. expected incidence (95% CI)	<i>P</i>
1 January 2020 to 30 June 2021*				
All patients	24.4 (23.6–25.2)	21.2 (20.5–21.9)	1.15 (1.10–1.20)	<0.001
Male patients	26.6 (25.5–27.8)	22.9 (21.9–24.0)	1.16 (1.10–1.23)	<0.001
Female patients	21.9 (20.9–23.0)	19.2 (18.2–20.2)	1.14 (1.07–1.21)	<0.001
Aged <6 years	18.6 (17.5–19.8)	15.1 (14.1–16.2)	1.23 (1.13–1.33)	<0.001
Aged 6–11 years	32.7 (31.2–34.3)	27.7 (26.3–29.2)	1.18 (1.11–1.26)	<0.001
Aged 12–17 years	23.5 (22.2–24.8)	22.2 (21.0–23.5)	1.06 (0.98–1.13)	0.13
Year 2020*				
All patients	23.7 (22.8–24.6)	20.9 (20.0–21.8)	1.13 (1.08–1.19)	<0.001
Male patients	25.7 (24.3–27.0)	22.6 (21.4–23.9)	1.13 (1.06–1.21)	<0.001
Female patients	21.5 (20.3–22.8)	18.9 (17.7–20.1)	1.14 (1.06–1.23)	<0.001
Aged <6 years	17.8 (16.5–19.2)	15.2 (14.0–16.5)	1.17 (1.06–1.30)	0.002
Aged 6–11 years	31.4 (29.6–33.4)	27.3 (25.6–29.1)	1.15 (1.07–1.24)	<0.001
Aged 12–17 years	23.4 (21.8–25.1)	21.5 (20.1–23.2)	1.09 (1.00–1.19)	0.06
First half of year 2021*				
All patients	25.8 (24.4–27.2)	21.8 (20.5–23.1)	1.18 (1.10–1.27)	<0.001
Male patients	28.6 (26.6–30.6)	23.5 (21.7–25.4)	1.22 (1.11–1.33)	<0.001
Female patients	22.8 (21.0–24.7)	19.8 (18.1–21.6)	1.15 (1.04–1.28)	0.008
Aged <6 years	20.2 (18.2–22.4)	15.0 (13.4–16.9)	1.34 (1.17–1.54)	<0.001
Aged 6–11 years	35.4 (32.7–38.3)	28.4 (26.0–31.1)	1.24 (1.12–1.38)	<0.001
Aged 12–17 years	23.6 (21.4–26.0)	23.6 (21.4–26.0)	1.00 (0.89–1.13)	0.96
Period with increased incidence of type 1 diabetes in 2020 (June 2020 to September 2020)†				
All patients	24.0 (22.0–26.1)	18.9 (17.2–20.8)	1.27 (1.13–1.43)	<0.001
Male patients	25.4 (22.7–28.5)	20.6 (18.1–23.4)	1.23 (1.05–1.44)	0.002
Female patients	22.4 (19.8–25.4)	16.9 (14.7–19.6)	1.33 (1.11–1.58)	<0.001
Aged <6 years	17.8 (15.1–21.0)	13.2 (10.8–16.0)	1.35 (1.07–1.71)	0.004
Aged 6–11 years	32.6 (28.7–37.1)	25.9 (22.4–29.8)	1.26 (1.06–1.50)	0.003
Aged 12–17 years	23.5 (20.2–27.3)	19.0 (16.1–22.5)	1.23 (1.00–1.51)	0.043
Period with increased incidence of type 1 diabetes in 2021 (March 2021 to June 2021)†				
All patients	25.5 (23.5–27.7)	20.2 (18.4–22.1)	1.27 (1.13–1.42)	<0.001
Male patients	28.8 (25.9–32.1)	21.7 (19.2–24.6)	1.33 (1.14–1.55)	<0.001
Female patients	22.1 (19.5–25.1)	18.4 (16.0–21.1)	1.20 (1.01–1.43)	0.023
Aged <6 years	20.6 (17.6–24.0)	14.1 (11.7–17.0)	1.46 (1.17–1.83)	<0.001
Aged 6–11 years	34.0 (30.0–38.5)	26.2 (22.8–30.2)	1.30 (1.09–1.54)	<0.001
Aged 12–17 years	23.4 (20.2–27.2)	21.6 (18.5–25.2)	1.09 (0.89–1.32)	1.00

Using Poisson regression models, expected incidences for 2020 were estimated based on the data from 2011 to 2019 and compared with observed incidences in 2020 by estimating IRRs with 95% CIs and two-sided *P* values. Incidences are presented as rate per 100,000 person-years with corresponding 95% CIs. Bold font indicates significance. *CIs of observed and expected incidences were adjusted for multiple inferences using the Bonferroni method separately for each stratum. †CIs of observed and expected incidences and CIs of IRRs were adjusted for multiple inferences using the Bonferroni method, and *P* values were adjusted for multiple testing using the Bonferroni-Holm method.

higher than the 231 cases in 2019, but significantly higher than 216 and 202 new cases in 2018 and 2017, respectively. However, the number of cases was too small and the time trend analysis too short to draw clear conclusions. Some other studies with fewer patients showed inconsistent results regarding a change in the incidence of pediatric type 1 diabetes during the pandemic (10,25–28). Supplementary Table 2 gives an overview of studies on incidence or number of cases

of pediatric new-onset type 1 diabetes during the COVID-19 pandemic compared with previous years.

A temporal association of the increased incidence of type 1 diabetes following the peak incidences of COVID-19 may be due to direct effects resulting from infections with SARS-CoV-2, but also to indirect effects resulting from the consequence of environmental changes caused by the pandemic itself or by pandemic containment measures.

In the case of direct effects, cytotoxic as well as immunologic effects could cause β -cell impairment. SARS-CoV-2 was suggested to cause diabetes through direct cytotoxicity of β -cells without autoimmunity (29). However, histopathologic studies showed conflicting evidence of whether SARS-CoV-2 directly affects pancreatic β -cells (30–33). In addition, this study found no increase in the frequency of autoantibody-negative type 1 diabetes in children and adolescents during those




Figure 3—IRRs of observed versus expected incidence of type 1 diabetes in children and adolescents from January 2020 to June 2021 by different pandemic-related periods. Error bars indicate 95% CIs. For period-specific analyses, statistical models were extended to include the period of diabetes onset and a term for an interaction of period at diabetes onset by year at diabetes onset. A Poisson regression model that included a binary variable indicating observed and predicted data and period as independent variables was used to compare the adjusted observed with the adjusted expected incidence of type 1 diabetes. Confidence limits of IRRs and *P* values were adjusted for multiple testing using the Bonferroni and Bonferroni-Holm methods, respectively.

periods, for which an increased observed versus expected incidence of type 1 diabetes has been demonstrated. This is consistent with a previous detailed analysis of the first half of 2020 (34). Therefore, the increase in the incidence of type 1 diabetes in children appears to be due to

immune-mediated type 1 diabetes. However, because autoimmunity and progressive β -cell destruction typically begin long before the clinical diagnosis of type 1 diabetes, we were surprised to see the incidence of type 1 diabetes followed the peak incidence of COVID-19 and also the

pandemic containment measures by only ~3 months.

Considering direct effects of COVID-19, infection with SARS-CoV-2 could trigger the development of new-onset type 1 diabetes in those with prediabetes with islet autoimmunity. Although infection with SARS-CoV-2 is typically very mild and often asymptomatic in children, a rare complication is multisystemic inflammatory syndrome (MIS-C) in children with COVID-19 (35). This hyper-inflammatory disorder presents 1–2 months after acute infection with SARS-CoV-2 with high fever, organ dysfunction, markedly elevated inflammatory markers, and formation of multiple autoantibodies. The progress from prediabetes to overt type 1 diabetes could be due to the cytokine storm seen in children after COVID-19, possibly as a different course of MIS-C with a focus on autoimmunity. However, immunologic findings seen in children with MIS-C and with new-onset type 1 diabetes differ (36,37). Unfortunately, sufficient data about the history of COVID-19 infections in children with new-onset type 1 diabetes are currently not available in the DPV. As mentioned above, no evidence of past SARS-CoV-2 infection was found in children with new-onset type 1 diabetes in the Finnish study (23). Furthermore, we found no association between the COVID-19




Figure 4—IRRs of observed versus expected incidence of type 1 diabetes in children and adolescents from January 2020 to June 2021 by month. IRRs estimated from trend models are presented as blue line with the corresponding 95% CI as gray area. For month-specific analyses, statistical models were extended to include the month of diabetes onset and a term for an interaction of period at diabetes onset by year at diabetes onset. A Poisson regression model that included a binary variable indicating observed and predicted data and month as independent variables was used to compare the adjusted observed with the adjusted expected incidence of type 1 diabetes. Confidence limits of IRRs and *P* values were adjusted for multiple testing using the Bonferroni and Bonferroni-Holm methods, respectively. IRRs were significantly elevated in June 2020 (*P* = 0.003), July 2020 (*P* < 0.001), March 2021 (*P* = 0.028), and June 2021 (*P* = 0.010).

incidences reported by the authorities and the increase in the observed over the expected incidence of type 1 diabetes in the following months. While numbers of COVID-19 cases were significantly higher in the second wave of the pandemic than in the first wave (1,2), the incidence of type 1 diabetes did not increase more after the second than after the first wave of the COVID-19 pandemic. Moreover, the seroprevalence for anti-SARS-CoV-2 antibodies in children and adolescents in Germany after the first wave of the pandemic was quite low, between 0.6 and 1.1% (38–40).

Considering indirect effects of COVID-19, the impact on type 1 diabetes incidence is not due to infection with SARS-CoV-2, but rather a consequence of environmental changes resulting from the pandemic itself or pandemic containment measures. Attention has been focused on the possibility that changes in lifestyle are a major factor in the rise of type 1 diabetes frequency (41). Because of social isolation, there was a drastic decrease in otherwise frequent and typical pediatric infectious diseases during the pandemic, such as viral respiratory and gastrointestinal tract infections (13,14). Common infections in early childhood have been shown to be a risk factor for the development of type 1 diabetes in children (5–9). This association would suggest a decrease in the incidence of type 1 diabetes. However, the opposite is the case; this study found an increase in the incidence of new-onset type 1 diabetes during the COVID-19 pandemic, despite the marked decrease in common pediatric infections.

It is important to note that infections could also protect against type 1 diabetes. According to the hygiene hypothesis, there is an inverse trend between the occurrence of infectious diseases in early life and the occurrence of autoimmune diseases (42). The biodiversity hypothesis, an extension of the hygiene hypothesis, states that decreased biodiversity in the external and internal exposure increases the risk of immune-mediated diseases (43). Therefore, the influence of environmental exposure on the risk of developing autoimmune diseases is of utmost importance (44). COVID-19 pandemic control efforts changed abruptly social contacts, the behavior of families with children, and child

health care practices, which may have resulted in a dramatic decrease in biodiversity in children, particularly in young children (45,46). This change in total environmental exposure, the exposome, could be the link to the increased incidence of type 1 diabetes in young children seen in our study (47,48).

In addition, increased stress from the pandemic itself or social isolation and school closure has been reported in children and adolescents, leading to poorer mental health and a high prevalence of anxiety and depressive symptoms (49–51). Among the environmental variables that play a role in the development of type 1 diabetes, psychological stress has been linked to the onset of the disease, and a wide range of studies point to the role of psychological stressors in the pathogenesis of type 1 diabetes (52–54).

A strength of our study is the population-based data from German children and adolescents. Furthermore, for comparisons with observed incidence, the estimates of the expected incidence in 2020 were derived from appropriate statistical methods, considering the observed increase in the incidence of type 1 diabetes over the past decade. Limitations of our study include that the multivariable Poisson regression model included only some potential confounders. Thus, residual confounding resulting from patient-, area-, and population-level confounders cannot be excluded. Potentially confounding factors include socioeconomic status, history of COVID-19, or family members with COVID-19. In addition, we can only speculate about underlying factors. The analysis relating incidence of type 1 diabetes and occurrence of COVID-19 is an ecologic analysis and thus prone to ecologic bias. However, an analysis of this association using individual-level data could not be performed, because the COVID-19 status of the patients was only available for a small proportion, and a suitable control group of children and adolescents without type 1 diabetes but known COVID-19 status was not available.

Further research is needed to understand the reasons for the increased incidence of type 1 diabetes following the waves of the pandemic and the role of infection with SARS-CoV-2 and also the measures taken to control the COVID-19 pandemic with its consequences, such as social isolation or psychological

stress. Additional longer-term studies on the incidence of type 1 diabetes during the pandemic are therefore essential.

In summary, this study found an increase in the incidence of type 1 diabetes in children and adolescents in Germany during the COVID-19 pandemic, with a time lag of ~ 3 months to the peak incidences of COVID-19 but also to the pandemic containment measures. The possible causes of this association are manifold and still unclear. However, our results suggest that indirect rather than direct effects of the pandemic are more likely to be the cause.

Acknowledgments. The authors thank A. Hungele and R. Ranz for support and development of the DPV documentation software (clinical data managers, Central Institute of Epidemiology and Medical Biometry [ZIBMT], Ulm University, Ulm, Germany) and Sascha Tittel for help with data analysis. The authors thank all centers participating in the DPV project (a list is available in the data supplement). 2020 data was presented at the ESPE (European Society for Paediatric Endocrinology) online congress 2021.

Funding. The DPV is supported through the German Federal Ministry for Education and Research within the German Centre for Diabetes Research (DZD) (82DZD14A02). Further financial support was received by the German Diabetes Foundation (DDS) (FP-0438-2021), the German Robert Koch Institute (RKI), and the German Diabetes Association (DDG).

The funding organization had no role in design or conduct of the study, collection, management, analysis, or interpretation of the data, preparation, review, or approval of the manuscript, or decision to submit the manuscript for publication.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. C.K. conceptualized the study, interpreted the analyses, wrote the initial manuscript, and reviewed and revised the manuscript. J.R. analyzed the data, designed and supervised the statistical analysis, and critically reviewed and revised the manuscript. A.J.E. analyzed the data and designed the analyses, contributed to the interpretation of results, and reviewed and revised the manuscript. K.S., H.B., D.K., M.S., S.H. and V.L. collected data, contributed intellectually to the research topics of the DPV initiative, and critically reviewed the scientific content of the manuscript. R.W.H. conceptualized the study, coordinated and supervised data collection, acquired funding for the study, and critically reviewed the manuscript for important intellectual content. All authors approved the final manuscript as submitted and agreed to be accountable for all aspects of the work. A.J.E. and R.W.H. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the

integrity of the data and the accuracy of the data analysis.

References

1. World Health Organization. Germany situation. Accessed 17 September 2021. Available at <https://covid19.who.int/region/euro/country/de>
2. Robert Koch Institut. Coronavirus SARS-CoV-2: Aktuelle Situationsberichte, Wochenberichte und COVID-19 Trends im Überblick. Accessed 17 September 2021. Available at https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Situationsberichte/Gesamt.html
3. Norris JM, Johnson RK, Stene LC. Type 1 diabetes-early life origins and changing epidemiology. *Lancet Diabetes Endocrinol* 2020; 8:226–238
4. Patterson CC, Harjutsalo V, Rosenbauer J, et al. Trends and cyclical variation in the incidence of childhood type 1 diabetes in 26 European centres in the 25 year period 1989–2013: a multicentre prospective registration study. *Diabetologia* 2019;62:408–417
5. Hyöty H. Viruses in type 1 diabetes. *Pediatr Diabetes* 2016;17(Suppl. 22):56–64
6. Stene LC, Oikarinen S, Hyöty H, et al. Enterovirus infection and progression from islet autoimmunity to type 1 diabetes: the Diabetes and Autoimmunity Study in the Young (DAISY). *Diabetes* 2010;59:3174–3180
7. Lönnerdal M, Lynch KF, Elding Larsson H, et al.; TEDDY Study Group. Respiratory infections are temporally associated with initiation of type 1 diabetes autoimmunity: the TEDDY study [published correction appears in *Diabetologia* 2018; 61:254]. *Diabetologia* 2017;60:1931–1940
8. Beyerlein A, Wehweck F, Ziegler AG, Pflueger M. Respiratory infections in early life and the development of islet autoimmunity in children at increased type 1 diabetes risk: evidence from the BABYDIET study [published correction appears in *JAMA Pediatr* 2017;171:94]. *JAMA Pediatr* 2013; 167:800–807
9. Ruiz PLD, Tapia G, Bakken IJ, et al. Pandemic influenza and subsequent risk of type 1 diabetes: a nationwide cohort study. *Diabetologia* 2018;61: 1996–2004
10. Unsworth R, Wallace S, Oliver NS, et al. New-onset type 1 diabetes in children during COVID-19: multicenter regional findings in the U.K. *Diabetes Care* 2020;43:e170–e171
11. Accili D. Can COVID-19 cause diabetes? *Nat Metab* 2021;3:123–125
12. Boddu SK, Aurangabadkar G, Kuchay MS. New onset diabetes, type 1 diabetes and COVID-19. *Diabetes Metab Syndr* 2020;14:2211–2217
13. Kruizinga MD, Peeters D, van Veen M, et al. The impact of lockdown on pediatric ED visits and hospital admissions during the COVID19 pandemic: a multicenter analysis and review of the literature. *Eur J Pediatr* 2021;180:2271–2279
14. Kuitunen I, Artama M, Mäkelä L, Backman K, Heiskanen-Kosma T, Renko M. Effect of social distancing due to the COVID-19 pandemic on the incidence of viral respiratory tract infections in children in Finland during early 2020. *Pediatr Infect Dis J* 2020;39:e423–e427
15. Tittel SR, Rosenbauer J, Kamrath C, et al.; DPV Initiative. Did the COVID-19 lockdown affect the incidence of pediatric type 1 diabetes in Germany? *Diabetes Care* 2020;43:e172–e173
16. Hofer SE, Schwandt A; Austrian/German DPV Initiative. Standardized documentation in pediatric diabetology: experience from Austria and Germany. *J Diabetes Sci Technol* 2016;10: 1042–1049
17. Robert Koch Institut. Coronavirus SARS-CoV-2: 7-Tage-Inzidenzen nach Bundesländern und Kreisen (fixierte Werte) sowie 7-Tage-Inzidenz der hospitalisierten COVID-19-Fälle nach Bundesländern (fixierte Werte), Stand: 27.12.2021. Accessed 24 September 2021. Available at https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Daten/Fallzahlen_Kum_Tab.html
18. German Federal Statistical Office. Population Table 12411-0013. Accessed 10 February 2021. Available at <https://www-genesis.destatis.de/genesis/online>
19. Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat Hum Behav* 2021;5:529–538. DOI: 10.1038/s41562-021-01079-8
20. Rosario AS, Kurth BM, Stolzenberg H, Ellert U, Neuhauser H. Body mass index percentiles for children and adolescents in Germany based on a nationally representative sample (KiGGS 2003–2006). *Eur J Clin Nutr* 2010;64:341–349
21. Skirvarhaug T, Stene LC, Drivvoll AK, Strøm H; Norwegian Childhood Diabetes Study Group. Incidence of type 1 diabetes in Norway among children aged 0–14 years between 1989 and 2012: has the incidence stopped rising? Results from the Norwegian Childhood Diabetes Registry. *Diabetologia* 2014;57:57–62
22. Rami-Merhar B, Hofer SE, Fröhlich-Reiterer E, Waldhofer T; Austrian Diabetes Incidence Study Group. Time trends in incidence of diabetes mellitus in Austrian children and adolescents <15 years (1989–2017). *Pediatr Diabetes* 2020;21: 720–726
23. Salmi H, Heinonen S, Hästbacka J, et al. New-onset type 1 diabetes in Finnish children during the COVID-19 pandemic. *Arch Dis Child* 27 May 2021 [Epub ahead of print]. DOI: 10.1136/archdischild-2020-321220
24. Mameli C, Scaramuzza A, Macedoni M, et al. Type 1 diabetes onset in Lombardy region, Italy, during the COVID-19 pandemic: the double-wave occurrence. *EClinicalMedicine* 2021;39:101067
25. Rabbone I, Schiaffini R, Cherubini V, Maffeis C; Diabetes Study Group of the Italian Society for Pediatric Endocrinology and Diabetes. Has COVID-19 delayed the diagnosis and worsened the presentation of type 1 diabetes in children? *Diabetes Care* 2020;43:2870–2872
26. Zubkiewicz-Kucharska A, Seifert M, Stępkowski M, Noczyńska A. Diagnosis of type 1 diabetes during the SARS-CoV-2 pandemic: does lockdown affect the incidence and clinical status of patients? *Adv Clin Exp Med* 2021;30:127–134
27. Lawrence C, Seckold R, Smart C, et al. Increased paediatric presentations of severe diabetic ketoacidosis in an Australian tertiary centre during the COVID-19 pandemic. *Diabet Med* 2021;38:e14417
28. Atlas G, Rodrigues F, Moshage Y, Welch J, White M, O’Connell MA. Presentation of paediatric type 1 diabetes in Melbourne, Australia during the initial stages of the COVID-19 pandemic. *J Paediatr Child Health* 2020;56:1654–1655
29. Hollstein T, Schulte DM, Schulz J, et al. Autoantibody-negative insulin-dependent diabetes mellitus after SARS-CoV-2 infection: a case report. *Nat Metab* 2020;2:1021–1024
30. Müller JA, Groß R, Conzelmann C, et al. SARS-CoV-2 infects and replicates in cells of the human endocrine and exocrine pancreas. *Nat Metab* 2021;3:149–165
31. Coate KC, Cha J, Shrestha S, et al.; HPAP Consortium. SARS-CoV-2 cell entry factors ACE2 and TMPRSS2 are expressed in the microvasculature and ducts of human pancreas but are not enriched in β cells. *Cell Metab* 2020;32:1028–1040.e4
32. Kusmartseva I, Wu W, Syed F, et al. Expression of SARS-CoV-2 entry factors in the pancreas of normal organ donors and individuals with COVID-19. *Cell Metab* 2020;32:1041–1051.e6
33. Fignani D, Licata G, Brusco N, et al. SARS-CoV-2 receptor angiotensin I-converting enzyme type 2 (ACE2) is expressed in human pancreatic β-cells and in the human pancreas microvasculature. *Front Endocrinol (Lausanne)* 2020;11:596898
34. Kamrath C, Rosenbauer J, Tittel SR, et al. Frequency of autoantibody-negative type 1 diabetes in children, adolescents, and young adults during the first wave of the COVID-19 pandemic in Germany. *Diabetes Care* 2021;44:1540–1546
35. Feldstein LR, Rose EB, Horwitz SM, et al.; Overcoming COVID-19 Investigators; CDC COVID-19 Response Team. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med* 2020;383:334–346
36. Consiglio CR, Cotugno N, Sardi F, et al.; CACTUS Study Team. The immunology of multisystem inflammatory syndrome in children with COVID-19. *Cell* 2020;183:968–981.e7
37. Ferreira RC, Simons HZ, Thompson WS, et al. IL-21 production by CD4+ effector T cells and frequency of circulating follicular helper T cells are increased in type 1 diabetes patients. *Diabetologia* 2015;58:781–790
38. Tönshoff B, Müller B, Elling R, et al. Prevalence of SARS-CoV-2 infection in children and their parents in southwest Germany. *JAMA Pediatr* 2021;175:586–593
39. Hippich M, Holthaus L, Assfalg R, et al. A public health antibody screening indicates a 6-fold higher SARS-CoV-2 exposure rate than reported cases in children. *Med (N Y)* 2021;2:149–163.e4
40. Kirsten C, Unrath M, Lück C, Dalpke AH, Berner R, Armann J. SARS-CoV-2 seroprevalence in students and teachers: a longitudinal study from May to October 2020 in German secondary schools. *BMJ Open* 2021;11:e049876
41. Egro FM. Why is type 1 diabetes increasing? *J Mol Endocrinol* 2013;51:R1–R13
42. Bach JF. Revisiting the hygiene hypothesis in the context of autoimmunity. *Front Immunol* 2021;11:615192
43. von Hertzen L, Hanski I, Haahela T. Natural immunity. Biodiversity loss and inflammatory diseases are two global megatrends that might be related. *EMBO Rep* 2011;12:1089–1093
44. Anaya JM, Ramirez-Santana C, Alzate MA, Molano-Gonzalez N, Rojas-Villarraga A. The autoimmune ecology. *Front Immunol* 2016;7:139
45. Finlay BB, Amato KR, Azad M, et al. The hygiene hypothesis, the COVID pandemic, and consequences for the human

- microbiome [published correction appears in *Proc Natl Acad Sci U S A* 2021;118:e2102333 118]. *Proc Natl Acad Sci USA* 2021;118:e2010217118
46. Renz H, Holt PG, Inouye M, Logan AC, Prescott SL, Sly PD. An exposome perspective: early-life events and immune development in a changing world. *J Allergy Clin Immunol* 2017;140:24–40
47. Traversi D, Rabbone I, Scaiola G, et al. Risk factors for type 1 diabetes, including environmental, behavioural and gut microbial factors: a case-control study. *Sci Rep* 2020;10:17566
48. Vatanen T, Franzosa EA, Schwager R, et al. The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature* 2018;562:589–594
49. Lee J. Mental health effects of school closures during COVID-19 [published correction appears in *Lancet Child Adolesc Health* 2020;4:e16]. *Lancet Child Adolesc Health* 2020;4:e16
50. Golberstein E, Wen H, Miller BF. Coronavirus disease 2019 (COVID-19) and mental health for children and adolescents. *JAMA Pediatr* 2020;174:819–820
51. Francisco R, Pedro M, Delvecchio E, et al. Psychological symptoms and behavioral changes in children and adolescents during the early phase of COVID-19 quarantine in three European countries. *Front Psychiatry* 2020;11:570164
52. Sipetic S, Vlajinac H, Marinkovi J, et al. Stressful life events and psychological dysfunctions before the onset of type 1 diabetes mellitus. *J Pediatr Endocrinol Metab* 2007;20:527–534
53. Thernlund GM, Dahlquist G, Hansson K, et al. Psychological stress and the onset of IDDM in children. *Diabetes Care* 1995;18:1323–1329
54. Nygren M, Carstensen J, Koch F, Ludvigsson J, Frostell A. Experience of a serious life event increases the risk for childhood type 1 diabetes: the ABIS population-based prospective cohort study. *Diabetologia* 2015;58:1188–1197